

$e^\pm$ Pair Cascades and Precursors in Gamma-Ray Bursts

P. Mészáros^{1,2,3}, E. Ramirez-Ruiz¹, M.J. Rees¹

¹Institute of Astronomy, University of Cambridge, Madingley Road, Cambridge CB3 0HA, U.K.

²Dpt. of Astronomy & Astrophysics, Pennsylvania State University, University Park, PA 16803

³Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106-4030

ABSTRACT

Gamma-ray burst sources with a high luminosity can produce electron-positron pair cascades in their environment as a result of back-scattering of a seed fraction of their original hard spectrum. The resulting spectral modifications offer the possibility of diagnosing not only the compactness parameter of the γ -ray emitting region but also the baryonic density of the environment external to the burst.

Subject headings: Gamma-rays: Bursts - X-rays - Cosmology: Miscellaneous

1. Introduction

The non-thermal γ -ray spectrum of Gamma-Ray Burst sources (GRBs) is thought to arise in shocks which develop beyond the radius at which the relativistic fireball from the initial event has become optically thin to scattering. However, the observed spectra are hard, with a significant fraction of the energy above the $\gamma\gamma \rightarrow e^\pm$ formation energy threshold, and a high compactness parameter can result in new pairs being formed outside the originally optically thin shocks responsible for the primary radiation. New pairs can be made as some of the initial energetic photons are backscattered and interact with other incoming photons. Such effects have been considered by Madau & Thompson (2000) and Thompson & Madau (2000), who investigated the acceleration of new pairs for a particular fireball model. Dermer & Böttcher (2000) considered the effect of pair formation for an external shock model of GRB, in which the compactness parameter is relatively low, while Madau, Blandford & Rees (2000) investigated the effects of Compton echos produced by pairs.

Here we present a simplified discussion of the generic effects of pair formation in a specific class of models. We suppose that the central engine gives rise to an unsteady baryonic wind, which is relativistic, carries a magnetic field, and lasts for a time t_w . A fraction of the wind energy is converted into γ -rays via internal shocks; the remaining wind energy drives a blast wave which decelerates as it sweeps up the external medium, and gives rise to the afterglow emission. The γ -rays would propagate ahead of the blast wave, leading to pair production (and an associated

deposition of momentum) into the external medium. The pair cascades saturate after the external (pair-enriched) medium reaches a critical bulk Lorentz factor, which is generally below that of the original relativistic wind. For external baryonic densities similar to those in molecular clouds the pairs can achieve scattering optical depths $\tau_{\pm} \lesssim 1$. Even for less extreme external densities the effect of the additional pairs can be substantial, increasing the radiative efficiency of the blast wave and leading to distortions of the original spectrum. This provides a potential tool for diagnosing the compactness parameter of the bursts and thus the radial distance at which shocks producing an observed luminosity can occur. It also provides a tool for diagnosing the baryonic density of the external environment of the bursts, and testing the association with star-forming regions.

2. Scattering and Two-photon Pair Formation

Consider an initial input radiation spectrum produced by the fireball of the form

$$\phi(\epsilon) = \frac{F}{m_e c^2 \epsilon_o^2 q} (\epsilon/\epsilon_o)^{-\beta} \text{ cm}^{-2} \text{ s}^{-1} \quad ; \quad \text{for } \epsilon \geq \epsilon_o, \quad (1)$$

where $F = L/4\pi d_L^2$ is the observed energy flux, $\epsilon = h\nu/m_e c^2$ is photon energy in electron rest-mass units, $\epsilon_o \sim 0.2 - 1.0$ is the break energy above which the photon number spectral index $\beta \sim 2 - 3$, and below which the spectral slope is flatter, e.g. $\alpha \sim -2/3$ for a simple low-energy cutoff synchrotron spectrum (although for the present purposes the exact low energy slope is unimportant), with $q \sim 1$ a normalization constant. For most values of the low and high energy slopes, the majority of the photons in the spectrum are near the break frequency.

For an $e^{\pm}$ moving away radially from the source of radiation with a velocity characterized by a Lorentz factor Γ , taking into account the Klein-Nishina drop-off in the scattering cross section above $\epsilon \sim 1$, the effective fraction of photons contributing to accelerating the electron is $\int_{\epsilon_o}^{\Gamma} \phi(\epsilon) d\epsilon = q^{-1} \epsilon_o^{\beta-2} (\beta-1)^{-1} [\Gamma^{-1} - \epsilon_o^{-1}] \simeq q^{-1} \epsilon_o^{-1} (\beta-1)^{-1}$ for $\beta > 1$. At a distance r in front of the radiation source, an $e^{\pm}$ can be accelerated to a maximum value of Γ satisfying

$$\frac{L}{4\pi r^2} \frac{1}{q \epsilon_o (\beta-1)} (\sigma_T/2) \Delta t \simeq \mu m_e c^2 \Gamma, \quad (2)$$

where the cross section near $\epsilon \sim 1$ is approximately $\sigma_T/2$, μm_e is the effective mass per scatterer (equation [3]), and Δt is the effective duration of the light pulse as seen by the electron. The latter is either $\Delta t \sim r/2c\Gamma^2$ (impulsive regime at small radii), or $\Delta t \simeq t_w$ (wind regime), depending on which one is smallest at a given radius, where t_w is the duration of the wind, or essentially the burst duration as seen by a distant observer. Equation (2) says that the total time-integrated momentum of the radiation intercepted by the electrons in the Thompson limit is converted into their kinetic energy of motion, taking into account the effective mean mass per scatterer. (This follows directly from the mean rate of momentum transfer per particle in a steady flow, e.g., as discussed in Madau & Thompson, 2000, their equation (2), (10) and (23), noting that our

definition of μ differs from theirs.) Here we define μ as the effective mass per scattering electron or positron in units of electron mass,

$$\mu m_e = \frac{\text{mass}}{\text{scatterer}} = m_p \frac{[1 + 2(n_{\pm}/n_p)(m_e/m_p)]}{[1 + (2n_{\pm}/n_p)]} \simeq \begin{cases} m_p & \text{for } n_{\pm} \ll n_p; \\ (2n_{\pm}/n_p)^{-1} m_p & \text{for } 1 < 2n_{\pm}/n_p < m_p/m_e; \\ m_e & \text{for } 2n_{\pm}/n_p > m_p/m_e. \end{cases} \quad (3)$$

Defining a reference radius

$$r_* = L\sigma_T/16\pi m_e c^3 = 5 \times 10^{19} L_{w50} \text{ cm} , \quad (4)$$

from equation (2) the maximum $e^{\pm}$ Lorentz factor is

$$\Gamma_{\pm}(r) \simeq \begin{cases} \mu^{-1/3} \left(\frac{1}{q\epsilon_o(\beta-1)} \right)^{1/3} (r_*/r)^{1/3} , & \text{for } \Delta t \sim (r/2c\Gamma^2); \\ \mu^{-1} \left(\frac{1}{q\epsilon_o(\beta-1)} \right) \left(\frac{2ct_w}{r_*} \right) (r_*/r)^2 , & \text{for } \Delta t \sim t_w. \end{cases} \quad (5)$$

The transition from the $\Gamma \propto r^{-1/3}$ to the steeper $\Gamma \propto r^{-2}$ occurs at a critical radius $r_c \ll r_*$, for which the Lorentz factor is Γ_c ,

$$\begin{aligned} r_c &= 2.2 \times 10^{14} \mu^{-2/5} (q\epsilon_o(\beta-1))^{-2/5} L_{w50}^{2/5} t_w^{3/5} \\ \Gamma_c &= 6.1 \times 10^1 \mu^{-1/5} (q\epsilon_o(\beta-1))^{2/15} L_{w50}^{1/5} t_w^{-1/5}. \end{aligned} \quad (6)$$

A second criterion for a maximum $e^{\pm}$ Lorentz factor comes from the pair formation threshold, since the incident photon ϵ and the back-scattered photon ϵ_r must satisfy $\epsilon\epsilon_r \geq 2$. The back-scattered photon, has, in the reference frame of the scattering $e^{\pm}$ moving with $\Gamma_{\pm}$, an energy of at most $\epsilon'_r \lesssim 1/2$; this photon cannot give rise to a further pair unless it collides with another photon with, in the lab frame, an energy exceeding $\sim 4\Gamma_{\pm}$. Thus the fraction of incident photons able to make pairs through the two-photon mechanism against target photons backscattered from $e^{\pm}$ moving with $\Gamma_{\pm}$ is $q^{-1}\epsilon_o^{-2} \int_{4\Gamma}^{\infty} \phi(\epsilon) d\epsilon = q^{-1}\epsilon_o^{\beta-2}(\beta-1)^{-1}(4\Gamma)^{1-\beta}$. The maximum Lorentz factor achievable by pairs before the two-photon cascade cuts off is that for which the compactness parameter has dropped to unity,

$$\ell(r) = \frac{L}{4\pi r^2} \frac{\epsilon_o^{\beta-2}}{q(\beta-1)} \frac{\sigma_T/3}{m_e c^2} \frac{1}{(4\Gamma_{\pm})^{\beta-1}} \Delta t \simeq 1 , \quad (7)$$

where the effective duration is as before. The maximum Lorentz factor for pair formation is then

$$\Gamma_{\pm}(r) \simeq \begin{cases} \left(\frac{2^{3-2\beta}}{3(\beta-1)} \right)^{1/(\beta+1)} \left(\frac{\epsilon_o}{q} \right)^{(\beta-2)/(\beta+1)} (r_*/r)^{1/(\beta+1)} , & \text{for } \Delta t \sim (r/2c\Gamma^2); \\ \left(\frac{4^{2-\beta}}{3(\beta-1)} \right)^{1/(\beta-1)} \left(\frac{\epsilon_o}{q} \right)^{(\beta-2)/(\beta-1)} \left(\frac{ct_w}{r_*} \right)^{1/(\beta-1)} (r_*/r)^{2/(\beta-1)} , & \text{for } \Delta t \sim t_w \end{cases} . \quad (8)$$

This yields a critical radius and Lorentz factor for the transition between the impulsive and wind dominated regimes of

$$\begin{aligned} r_c &= \left(\frac{2^{7-3\beta}}{9(\beta-1)^2} \right)^{\frac{1}{\beta+3}} \left(\frac{\epsilon_o}{q} \right)^{\frac{2\beta-4}{\beta+3}} r_* \left(\frac{ct_w}{r_*} \right)^{\frac{\beta+1}{\beta+3}} \simeq 5 \times 10^{14} L_{w50}^{2/5} t_{w1}^{3/5} , \\ \Gamma_c &= \left(\frac{2^{-2(\beta-1)}}{3(\beta-1)} \right)^{\frac{1}{\beta+3}} \left(\frac{\epsilon_o}{q} \right)^{\frac{\beta-2}{\beta+3}} \left(\frac{r_*}{ct_w} \right)^{\frac{1}{\beta+3}} \simeq 3 \times 10^1 L_{w50}^{1/5} t_{w1}^{-1/5} , \end{aligned} \quad (9)$$

where the numerical values in the second equation of both lines are calculated for $\beta = 2$, $\epsilon_o \simeq q \simeq 1$, $L = 10^{50} L_{w50} \text{ erg s}^{-1}$ and $t_w = 10 t_{w1} \text{ s}$. (At early times $t < t_w$ the radius r_c is smaller, $\propto t^{3/5}$, since it takes t_w for the entire photon energy to build up). The maximum pair Lorentz factor is shown in Figure 1 for nominal parameter values. The pair formation limit (9) is somewhat more restrictive than the scattering limit (6). Also, unlike the scattering limit, the pair formation limit depends exponentially on the photon number slope β . However, for the canonical value $\beta = 2$ the two power law dependences are the same and the numerical values are close to each other. In what follows we shall use the pair formation limit of equations (8) and (9).

3. Pair-precursor and Kinematics

Given a certain external baryon density n_p at a radius r outside the shocks producing the GRB primordial spectrum, the initial Thomson scattering optical depth is $\tau_i \sim n_p \sigma_T r$ and a fraction τ_i of the primordial photons will be scattered back, initiating a pair cascade. Since the photon flux drops as r^{-2} , for a uniform (or decreasing) external ion density most of the scattering occurs between r and $r/2$, and the scattering and pair formation may be approximated as a local phenomenon. This pair-dominated plasma, as long as its density $n_{\pm} \lesssim n_p (m_p/2m_e)$, is initially held back by the inertia of its constituent ions component, provided that the pairs remain coupled to the baryons. (The latter is likely to be the case in the presence of even weak magnetic fields, e.g. Madau & Thompson 2000, or plasma wave scattering, e.g. Lightman, 1982. Moreover, the initial magnetic field strength can also increase as consequence of instabilities caused by an initial pair streaming relative to ions. In what follows, we assume that this coupling is effective).

The pairs (together with the ions) start acquiring a significant velocity $\Gamma_{\pm} \gtrsim 1$ only after their density $n_{\pm} \gtrsim n_p (m_p/2m_e)$, when the effective mass per scatterer $\mu m_e \sim m_e$. After the pairs start being accelerated, the pair cascade can continue multiplying as long as the compactness parameter of equation (8) is $\ell \gtrsim 1$. Both the compactness ℓ and the maximum $\Gamma_{\pm}$ decrease for increasing radius according to equations (7) and (8), and the cascade process shuts off for $\ell \lesssim 1$ at a radius where the maximum $\Gamma_{\pm} \sim 1$,

$$r_{\ell} \sim (4r_* c t_w / 3)^{1/2} \sim 4 \times 10^{15} L_{w50}^{1/2} t_{w1}^{1/2} \text{ cm}. \quad (10)$$

On the other hand, the blast wave producing the shock(s) responsible for the afterglow spectrum starts decelerating (i.e. its initial Lorentz factor $\Gamma_f \sim \eta = 10^2 \eta_2$ starts to decrease as a negative power law in radius) at the deceleration radius

$$r_d \simeq (3L t_w / 4\pi n_p m_p c^2 \eta^2)^{1/3} \simeq 3 \times 10^{15} L_{w50}^{1/3} t_{w1}^{1/3} \eta_2^{-2/3} n_3^{-1/3} \text{ cm}, \quad (11)$$

where we normalized to an external baryon density $n_p = 10^3 n_3 \text{ cm}^{-3}$. (This expression is valid provided that the deceleration radius exceeds $c t_w \eta^2$. Otherwise the deceleration starts before the outflow is over, and t_w should be replaced by t in the expression). A larger value $n_p \geq 10^3 n_3 \text{ cm}^{-3}$ would give $r_d \lesssim r_{\ell}$. (Such a higher density would, e.g., be required for pair effects to be important

in an external shock model, where the γ -rays arise at r_d). In an internal shock model the primary γ -rays themselves arise at a smaller radius

$$r_i \sim ct_v \eta^2 \sim 3 \times 10^{13} t_{v-1} \eta_2^2 \text{ cm} \quad (12)$$

where $t_v = 10^{-1} t_{v-1} \gtrsim 10^{-3}$ s is the variability timescale in the outflow. The criterion for an $e^\pm$ cascade to form ahead of the fireball producing a primary spectrum (1) is that r_i (or r_d for an external shock model) is smaller than r_ℓ . Pair formation occurs then for $r \leq r_\ell$.

The minimum number of pairs is formed just inside r_ℓ , where they are created essentially at rest, and their density can reach $n_\pm \sim (m_p/2m_e)n_p \sim 10^3 n_p$, at which point the mass per scatterer has become comparable to the electron mass. This represents an increase in the electron (or positron) scattering opacity by a multiplication factor of $k_p \sim 2 \times 10^3$. As the density exceeds this value at r_ℓ the pairs are pushed by scattering beyond r_ℓ , where the condition $\ell < 1$ prevents further multiplication.

For radii $r < r_\ell$, and as long as the blast wave has not reached that radius, pair formation exceeding the above value $n_\pm \sim 10^3 n_p$ can continue to occur, with the pairs being accelerated by scattering until they reach the maximum Lorentz factor (8). Pair production will be most copious at smaller radii $r_i < r < r_c$, where the pairs are accelerated to a maximum Lorentz factor $\Gamma_\pm(r) \lesssim \Gamma_c(r/r_c)^{-1/3}$. At each step of the cascade the Lorentz factor of the new generation of pairs approximately doubles, and the maximum pair multiplication factor of the cascade can be estimated from the number s of pair generations required to go from $\Gamma \sim 1$ to $\Gamma_\pm(r)$ given by equation (8). At r_c this is $s(r_c) \sim \log \Gamma_c / \log 2 \sim 5$, and for $r_i \lesssim r$ it is limited to $s(r_i) \lesssim 7.5$. This represents an extra multiplication factor $k_a \sim 2^s \sim \Gamma_c(r/r_c)^{-1/3} \sim 30 - 170$ in the electron opacity for $r_i \leq r \leq r_c$, in addition to the previous factor $k_p \sim 2 \times 10^3$ achieved before acceleration starts. The critical value is

$$k_c = k_p k_a(r_c) \sim (m_p/m_e) \Gamma_c \sim 5 \times 10^4 L_{w50}^{1/5} t_{w1}^{-1/5}, \quad (13)$$

with $k(r) = k_c(r/r_c)^{-1/3}$ for $r_i \leq r_c$, while $k(r) \sim k_c(r/r_c)^{-2}$ for $r_c \leq r \leq r_\ell$ drops to m_p/m_e at r_ℓ , and to zero beyond that. In a more detailed calculation, the number of cascades depends somewhat on the radiation spectral index β , here set equal to 2. For a steeper spectrum, the number of cascades is reduced (e.g. Ramirez-Ruiz *et al.* 2000). The maximum pair optical depth is achieved at r_c ,

$$\tau_{\pm c} \sim k_c n_p \sigma_T r_c = \min[2 \times 10^{-5} n_p L_{w50}^{3/5} t_{w1}^{2/5}, 1]. \quad (14)$$

The pair opacity scales as $\tau_\pm \propto r^{2/3}$ for $r_i \lesssim r \lesssim r_c$ and $\tau_\pm \propto r^{-1}$ for $r_c \lesssim r \lesssim r_\ell$. For an external density greater than

$$n_{p,c} \simeq 10^5 L_{w50}^{-3/5} t_{w1}^{-2/5} \text{ cm}^{-3}, \quad (15)$$

the pair density inside r_c is prevented from multiplying beyond a value corresponding to $\tau_\pm \sim 1$, due to self-shielding. The number of pairs created at r_c is

$$N_{\pm c} \sim n_p k_c (4\pi/3) r_c^3 \sim 3 \times 10^{49} n_p L_{w50}^{7/5} t_{w1}^{8/5}, \quad (16)$$

which scales as $N_{\pm} \propto r^{8/3}$, r for r below or above r_c . The maximum value for $n_p = n_{p,c}$ is $N_{\pm,c,max} \sim 3 \times 10^{54} L_{w50}^{4/5} t_{w1}^{6/5}$ at $r = r_c$, and $N_{\pm,\ell,max} \sim 3 \times 10^{55} L_{w50}^{9/10} t_{w1}^{11/10}$ at r_{ℓ} . The pair cloud extends over a range of radii comparable to r , much larger than the photon pulse radial width ct_w .

Figure 2 shows the schematic world-lines of three blast waves expanding into an external medium. At each radius $r_i < r < r_{\ell}$ pairs form ahead of r_i . While for $r \sim r_{\ell}$ the pairs are sub-relativistic and do not propagate significantly, for smaller radii they move relativistically with an initial speed $\Gamma_{\pm}$, with $\Gamma_{\pm} < \eta$ in general, although at very small radii, and for times $t < t_w$, $\Gamma_{\pm} > \eta$ may be possible. If the blast wave did not exist, then the pair-enriched material (moving out with a Lorentz factor that is larger at smaller r) would pile up at a radius r_l after a time r_l/c . However, the pair-rich external medium would generally carry less energy and inertia than the relativistic wind itself; it therefore starts being decelerated by the external medium at a smaller radius than a blast wave with the same Lorentz factor would be, so that the latter always overtakes it and sweeps up the pair-enriched external medium. The resultant outer shock is more likely to be radiative, but may be weaker, because the ambient material might already be moving outward relativistically before the blast wave hits it.

For a relatively low external density such as $n_p = 10^3 \text{cm}^{-3}$, as in curve (a) of Figure 2, the radius $r_c < r_d$, and the pairs are swept up by a fireball expanding at its original speed $\eta \sim \text{constant}$, and deceleration of the fireball begins sometime after the pairs have been swept up. In this case the main afterglow, produced by the deceleration of the pair-enriched fireball will be characterized by a larger emission measure than in the usual case, due to the extra pairs in the swept-up matter, but will not otherwise exhibit any irregularities in its time development. For a larger external density such as 10^6cm^{-3} , corresponding to curve (b) of Figure 2, the radius $r_c \sim r_d$ and the main part of the afterglow, caused by the deceleration of the fireball baryons and the pairs formed at early stages, continues to sweep up further pairs which were formed at or beyond the deceleration radius (in curve (c), with low η , deceleration occurs at radii $r > r_{\ell}$). These new pairs change the emission measure of the deceleration shock, since their optical depth $\tau_{\pm} \sim 1$ is larger than that of the swept up baryonic electrons. If $\tau_{\pm} \sim 1$ is reached, the original spectrum may be modified by Comptonization, and may approach a quasi-thermal shape. We note that the optical depth cannot substantially exceed unity. This is because high-energy non-thermal photons would be comptonised by pairs, and after a few scatterings would be degraded below the pair-production threshold. So, however intense the radiation incident on its inner edge, a shell of pair-dominated plasma cannot build up an opacity much larger than unity. It will come to a quasi-steady state in which the luminosity from its outer surface (comptonised synchrotron radiation and annihilation radiation) equals the non-thermal radiation shining on its inner surface.

4. Observable Pair-precursor Effects

The energy in the accelerated pairs is $E_{\pm} \propto N_{\pm} \Gamma_{\pm} r^3 \propto r^{7/3}$ for $r \lesssim r_c$ and $E_{\pm} \propto r^{-1}$ for $r_c \lesssim r \lesssim r_{\ell}$. The maximum number of pairs created is

$$E_{\pm c} \simeq N_{\pm} m_e c^2 \Gamma_c = \min[10^{45} n_p L_{w50} t_{w1}, 10^{50} L_{w50}^{2/5} t_{w1}^{3/5}] \text{ erg}, \quad (17)$$

where the second number is the limit $N_{\pm, mx}$ obtained for $n_p = n_{pc}$, at which the saturation value of $\tau_{\pm} \sim 1$ is obtained. The pair energy never exceeds more than a fraction of the initial wind energy, so the deceleration radius (equation [11]) is unaffected, while the pairs which are produced and accelerated ahead of the blast wave, are decelerated at a (smaller) radius

$$r_{d\pm} \sim 2.5 \times 10^{15} E_{\pm 50}^{1/3} n_3^{-1/3} (\Gamma_{\pm}/30)^{-2/3} \text{ cm}, \quad (18)$$

before the ejecta itself starts to decelerate.

The total number of protons in the ejecta wind is $N_p = E/\eta m_p c^2 = (2/3) \times 10^{52} L_{w50} t_{w1} \eta_2^{-1}$. The number of protons in the (pair enriched) external medium which are swept up by the time the blast wave reaches its deceleration radius r_d is $N_{ps} = N_p \eta^{-1} = (2/3) \times 10^{50} L_{w50} t_{w1} \eta_2^{-2}$. The maximum number of swept-up pairs can exceed the number of baryons by a large factor. However the inertia of the pairs is less than that of the baryonic ejecta, since $(N_{\pm, c, mx}/N_p) \sim 5 \times 10^2 L_{w50}^{-1/5} t_{w1}^{1/5} \eta_2 \lesssim (m_p/m_e)(\eta/\Gamma_c) = 6 \times 10^3 L_{w50}^{-1/5} t_{w1}^{1/5} \eta_2$ at r_c , and $(N_{\pm, \ell, mx}/N_p) \sim 5 \times 10^3 L_{w50}^{-1/10} t_{w1}^{1/10} \eta_2 \leq (m_p/m_e)\eta = 2 \times 10^5 \eta_2$ at r_{ℓ} .

The pairs will modify the usual properties of the deceleration shocks and the afterglow emission. The total energy available in the afterglow is not changed: it is still, essentially, the kinetic energy of the relativistic wind, minus the fraction dissipated (and converted into prompt gamma rays) in internal shocks. But the lepton/proton ratio in the ejecta can be much larger than usual. This increases the radiative efficiency significantly, since most of the particles are $e^{\pm}$. The resultant radiation will thus be softer than in the usual picture, because the same energy density has to be shared among a larger number of particles ($N_{\pm}/N_p \gg 1$). In the reverse shock that occurs after the baryonic+pair ejecta starts being decelerated by the external medium, the comoving frame peak random electron Lorentz factor is $\gamma_{\pm, m, r} \sim (E\eta^{-1}/N_{\pm} m_e c^2)$. Even for $\tau_{\pm}$ substantially less than unity, e.g. for an external density $n = 10^3 n_3 \text{ cm}^{-3}$ (below the critical value $n_{p, c}$) for which the deceleration radius $r_d \sim r_c$, the reverse shock random lepton Lorentz factor is $\gamma_{\pm, m, r} \sim 30 E_{w51}^{-1/5} t_{w1}^{-3/5} n_3^{-1} \eta_2^{-1}$. The pairs should not affect, however, the random turbulent magnetic field strength, which is in pressure equilibrium with the forward shock at some fraction ϵ_B of the equipartition value, $B' \sim (\epsilon_B 8\pi n m_p c^2)^{1/2} \eta \sim 6 \times \epsilon_B^{1/2} n_3^{1/2} \eta_2 \text{ G}$. Thus the observed reverse synchrotron peak frequency

$$\nu_{\pm, sy, r} \sim 10^6 B' \gamma^2 \eta (1+z)^{-1} \sim 6 \times 10^{13} E_{w51}^{-2/5} t_{w1}^{-6/5} \epsilon_B^{1/2} n_3^{-3/2} \eta_2^{-1} (1+z)^{-1} \text{ Hz} \quad (19)$$

would be in the far IR, as opposed to the optical/UV of the usual baryon-dominated prompt reverse flash.

The forward shock afterglow radiation, normally in the γ /X-ray range, would be unaffected outside r_ℓ . However, up to about the time when deceleration starts, there will be a “pair pick-up” photon pulse, when the protons and electrons of the relativistic wind moving with $\eta > \Gamma_c$ sweep up the pair-enriched external medium. The protons are not decelerated by the pairs, so there is only a subsonic reverse compression wave in the proton ejecta, but there will be a mildly relativistic forward shock in the picked-up pairs. The $e^\pm$ random Lorentz factor will be equal to the bulk kinetic energy it has in the proton frame, e.g. $\gamma_{\pm,m,f} \sim \eta/\Gamma_\pm \sim 3L_{w50}^{-1/5} t_{w1}^{1/5} \eta_2$ near r_c . The comoving random magnetic field in the shocked pair fluid is $B' \sim (8\pi\epsilon_B N_{\pm,c} m_e c^2 / 4r_c^3)^{1/2} (\eta/\Gamma_\pm) \sim 20\epsilon_B^{1/2} L_{w50}^{1/5} t_{w1}^{1/10} n_3^{1/2} \eta_2$ G for $n_p = 10^3 n_3 \text{cm}^{-3}$, corresponding to a pair-pickup pulse synchrotron peak frequency in the observer frame of

$$\nu_{\pm,sy,f} \sim 2 \times 10^{10} \epsilon_B^{1/2} L_{w50}^{-1/5} t_{w1}^{5/10} n_3^{1/2} \eta_2^3 (1+z)^{-1} \text{ Hz}, \quad (20)$$

and a synchrotron power law above this frequency.

There could also be additional pair-precursor signatures which are not associated with the ejecta blast wave, but with the dynamics of the pre-accelerated pair-enriched plasma. The r -dependent $\Gamma_\pm$ of the pair-enriched external plasma will lead to internal shocks. Pair regions at $r < r_c$ whose $\Gamma_\pm$ differ by order unity will collide at radii $\propto r\Gamma_\pm^2 \propto r^{1/3}$, so for sufficiently short variability times these shocks would occur between r_i and r_c , i.e. up to an observer time $t_c \sim r_c/c\Gamma_c^2 \sim 15t_{w1}$ s, independent of the wind luminosity L . (For $r_c < r < r_\ell$ the collision radii are $\propto r^{-3}$, so the shocks would tend to pile up at r_ℓ at an observer timescale $t_\ell \sim r_\ell/c \sim 1$ day, provided the main part of the fireball wind has not caught up with it before. This could be the case for a jet-like wind for pairs in a rim of angular width Γ_c^{-1} around the jet). The pair internal shocks at $r \sim r_c$ can produce a Fermi accelerated power law $e^\pm$ spectrum above $\gamma_{\pm m} \sim 1$ and a turbulent comoving field $B'_\pm \lesssim 2 \times 10^2$ G, leading to a synchrotron spectrum whose observer-frame peak frequency $\nu_{\pm m} \sim 6 \times 10^9$ Hz would be self-absorbed, with a power law extension above it which would be optically thin at higher frequencies.

The above observational signatures would be present even if $\tau_\pm$ is low, as expected for external ion densities $n_p \lesssim n_{p,c} \sim 10^5 L_{w50}^{-3/5} t_{w1}^{-2/5} \text{cm}^{-3}$ around the burst. An additional effect of interest, for external densities in excess of this which lead to a pair screen of optical depth $\tau_\pm \sim 1$, is that a quasi-thermal pulse of X-rays could accompany the burst, caused by upscattering of diffuse progenitor stellar photons. A collapsing massive progenitor leading to a GRB is likely to be highly super-Eddington for sometime after the GRB event, e.g. $L_* \sim 10^4 L_{Ed} \sim 10^{43} L_{*43} \text{ erg/s}$. The density of photons of energy $\epsilon_* \sim 10$ eV near r_c , which would be quasi-isotropized due to the condition $\tau_\pm \sim 1$ at r_c (and beyond, where $\Gamma_\pm \rightarrow 1$ at r_ℓ), would be $n_* \sim 10^{13} L_{*43} \epsilon_{*10}^{-1} L_{w50}^{-4/5} t_{w1}^{-6/5} \text{cm}^{-3}$ near r_c . The screen with $\tau_\pm \sim 1$ moving with $\Gamma_c \sim 30$ sweeps up a total number of photons $N_* \sim 3 \times 10^{57} L_{*43} \epsilon_{*10}^{-1} L_{w50}^{2/5} t_{w1}^{3/5}$. The mean energy per photon and the time-integrated total energy of the upscattered precursor is

$$\begin{aligned} \epsilon_X &\sim 10 \epsilon_{*10} L_{w50}^{2/5} t_{w1}^{-2/5} (1+z)^{-1} \text{ keV} \\ E_X &\sim 5 \times 10^{49} L_{*43} \epsilon_{*10}^{-1} L_{w50}^{4/5} t_{w1}^{1/5} (1+z)^{-1} \text{ erg.} \end{aligned} \quad (21)$$

This would last until the pairs are swept up by the ejecta, $t_X \lesssim t_c \sim r_c/c\Gamma_c^2 \sim 15t_{w1}$ s.

5. Discussion

We have discussed, in the context of a standard internal/external shock model of gamma-ray bursts (which are normally assumed to occur after the original fireball has become optically thin) the $\gamma\gamma \rightarrow e^\pm$ cascades triggered by the back-scattering of seed gamma-ray photons on the external medium. This effect can modify the initial scattering optical depth of the outflow at radii $r \lesssim r_\ell \simeq 4 \times 10^{15} L_{w50}^{1/2} t_{w1}^{1/2}$ cm (equation [10]), which is comparable to the radii of external shocks of equation (11) at which the afterglow begins, and is generally larger than the typical internal shock radii given by equation (12). The spectral effects of the pairs on the burst and the afterglow can be substantial, and within radii $\sim r_l$ they can affect the dynamics.

Pair production can increase the optical depth outside of the shocks by up to $\lesssim 10^5$; self-shielding ensures that the maximum scattering optical depths achievable by the pairs is $\tau_\pm \sim 1$. For typical interstellar densities the pair opacity $\tau_\pm \ll 1$, which does not significantly affect the gamma-ray spectrum. The number of pairs may nonetheless be large enough to increase the radiative efficiency and soften significantly the radiation spectrum of the afterglow reverse shock, where the same energy is shared among a number of $e^\pm$ which can exceed that of the original e^- and p^+ of the ejecta.

The pair production processes themselves (determining r_c , r_l , equations [9], [10]) just depend on the “seed” γ -ray photon flux (which are here postulated to come from internal shocks). The manifestations depend on the external density and on the initial dimensionless entropy or bulk Lorentz factor η . The external baryon density n_{ext} determines the optical depth that can be built up through back-scattering and pair multiplication. This affects whether the pair optical depth gets up to unity, with smearing and reprocessing of the primordial γ -ray spectrum, or whether it merely makes the blast wave more radiative. Madau and Thompson (2000) have made this point, in the context of a specific fireball model.

The external density (along with the initial Lorentz factor η) determines when the outer shock and the reverse shock become important and whether this happens within the radius already polluted with pairs (and pre-accelerated by radiation pressure before the shock hits). There are two rather different cases depending on whether or not η^2 is less than r_l/ct_w . In the former case the external shock responsible for the afterglow occurs beyond the region “polluted” by new pairs, while in the second case the afterglow shock may experience, after starting out in the canonical manner, a “resurgence” or second kick as its radiative efficiency is boosted by running into an $e^\pm$ -enriched gas. Internal shocks in the pair-dominated external plasma can lead to self-absorbed radiation at $\sim 10^9 - 10^{10}$ Hz, while the swept-up pairs can also contribute a $10^{11} - 10^{12}$ Hz ‘prompt’ signal, which precedes the onset of the standard deceleration afterglow phase.

Additional effects are expected when $\tau_\pm \rightarrow 1$. This requires external baryon densities at radii

$r < r_\ell$ of $n_p \gtrsim n_{c,p} \sim 10^5 L_{w50}^{-2/5} t_{w1}^{-3/5} \text{ cm}^{-3}$. Such high densities would only be expected if the burst is associated with a massive star in which prior mass loss led to a dense circumstellar envelope. The pair optical depth saturates to $\tau_\pm \sim 1$ and in addition to an increased efficiency and softer spectrum of the afterglow reverse shock, the original gamma-ray spectrum of the GRB will be modified as well. The specific nature of this spectral modification depends on the value of the luminosity, which influences (equation [9]) the bulk Lorentz factor of the reprocessing pair cloud before it has been swept up by the ejecta, and also on the extent to which the outflow is beamed. One of the consequences of such a critical external density leading to $\tau_\pm \sim 1$ would be the presence of an X-ray quasi-thermal pulse, whose total energy may be a few percent of the total burst energy. In the case of even more extreme densities, there are other interesting possibilities. For instance, dense blobs of Fe-enriched thermal plasma would emit strong recombination features, as well as annihilation radiation features, if the normal electron density were augmented by extra pairs.

Even in isotropic situations, the spectrum would be modified by transmission through a pair plasma of optical depth unity. The effect is maximal for photons of energy $m_e c^2$ in the frame of the pair plasma: for higher energies, the Klein-Nishina cross-section is smaller; for lower energies, the scatterings are almost elastic. If, for instance, $\Gamma_c \sim 40$, the gamma-ray spectrum around photon energies $\epsilon \sim \Gamma_c m_e c^2 \sim 20 \text{ MeV}$ would be depressed by a factor ~ 1 relative to its original value, smoothly rejoining its original value above and below that energy. For a beamed primary output, however, there would be a suppression at lower energies (where the scattering is in the Thompson regime) because the scattered photons would be spread over a wider angle. Pair-induced processes would therefore yield evidence on the beaming properties of the bursts.

Irrespective of the external density, the processes discussed here suggest that bursts and afterglows may have a more complex spectrum and time-structure than ‘standard’ models suggest. But the effects are especially interesting when the external density is high: they probe the environment of GRBs, and thus can offer clues to the nature of the progenitor stars, and their location within the host galaxy. For instance, a quasi-thermal X-ray pulse accompanying the gamma-ray emission could be indicative of an external circum-burst density of at least $10^4 - 10^5 \text{ cm}^{-3}$. While quasi-thermal X-ray pulses might also arise due to other reasons, e.g. from an underlying optically thick central engine, if the X-ray luminosity scales as equation (21) and is accompanied by radio or far-IR signals such as in equations (19)(20), this could be indicative of birth in a dense environment from a massive progenitor.

Detailed Monte Carlo simulations (Ramirez-Ruiz *et al.*, 2001) should provide a more detailed assessment of the self-consistent spectrum of a GRB in the presence of self-induced pair formation.

This research has been supported by NASA NAG5-9192, the Guggenheim Foundation, the Sackler Foundation, NSF PHY94-07194 and the Royal Society. We are grateful to R. D. Blandford, C. Dermer, P. Madau, C. Thompson and the referee for useful comments.

REFERENCES

- Dermer, C & Böttcher, M., 2000 ApJ 534, L155
 Lightman, A.P, 1982, ApJ 253, 842.
 Madau, P & Thompson, C, 2000 ApJ, 534, 239
 Madau, P, Blandford, R & Rees, M.J., 2000, ApJ, 541, 712.
 Ramirez-Ruiz, E, *et al.*, 2001, in preparation
 Thompson, C & Madau, P, 2000 ApJ, 538, 105




Fig. 1.— Schematic plot of the maximum $e^\pm$ bulk Lorentz factor $\Gamma_\pm$ as a function of radius.




Fig. 2.— This diagram shows, for three illustrative cases, how the burst would be affected by pair-enrichment. The axes (logarithmic) are r versus $t - (r/c)$, where t is time measured by a distant observer, and is zero when the burst is observed to start. (In this plot, light rays are horizontal lines). The primary gamma-ray emission is assumed to continue, with a quasi-steady luminosity L , for a time t_w (as would happen if, for instance, it came from internal shocks induced by wind variability on time-scales $\ll t_w$). The criterion for runaway pair production would be satisfied within the ambient material out to a radius r_ℓ . (cf equation [10]; t replaces t_w in this equation for times $t < t_w$.) The associated absorption of momentum would itself accelerate this pair-enriched material to an r -dependent Lorentz factor (see Figure 1). The wind generally carries more momentum than the gamma rays, and drives a relativistic blast wave that eventually sweeps up all the pair-enriched medium. Three illustrative cases are depicted. In case (a), the external medium has a low density, and the blast wave, with high η , sweeps up all the pair-enriched medium before it has been much decelerated: the effects of the pairs are then observed primarily during the burst itself. In case (b), with higher external density, deceleration occurs at radii $< r_\ell$, and the blast wave is still moving through pair-enriched material during the afterglow. When the ambient medium is dense, the pairs may provide an optical depth of unity, so the primary burst itself would be reprocessed, and its short time-structure smeared out. Case (c) corresponds to a lower η . The sweeping-up of pairs then occurs during the afterglow (modifying the radiative efficiency of the outer shock) even if the external density is low and there has not (as in case b) already been deceleration.